

Outrageous Amounts of Mercury Being Injected Into Children

News from Mariandina Research Foundation



This article may be of interest to you as I am sure we are all concerned as to what our children are being injected with, as responsible parents and grand-parents, there is a NEED TO KNOW. Rather frightening don't you think? No wonder the Prime Minister of England refused to have his child immunised.

One hundred years ago, children received 1 vaccine (the smallpox vaccine). Forty years ago, children received 5 vaccines routinely (diphtheria, pertussis, tetanus, polio, and smallpox vaccines) and as many as 8 shots by 2 years of age.

Children now receive 52 vaccines, in the form of 15 shots, by the time they are 6 months of age they receive all the recommended shots, including the Prevnar pediatric pneumonia shot.

Vaccines contain contain THIMERSOL (mercury), MSG, aluminum, formaldehyde, sucrose and phenoxyethanol, which is antifreeze, among many other things.

Thimersol, a vaccine ingredient, is nearly 50% mercury.

Mercury is a NEUROTOXIN.

EPA 'safe' levels are: 1 microgram per 1.0 kilogram of body weight per day.

Vaccines contain 12.5 to 25.0 micrograms of mercury, and a 'well baby' visit can see your child have between 50 and 62.5 mcgs of MERCURY injected into their bloodstream.

The CDC (US) has found a trend linking autism to mercury laden vaccines.

Thimerosal is a registered pesticide with the Department of Pesticide Registration of the Environmental Protection Agency.

Vaccines given:

Day of Birth: Hepatitis B contains 12 mcg mercury which is 30 times above the 'safe' level.

At 4 month: DTaP and HiB on same day. These contain 50 mcg mercury which is 60 times above the 'safe' level.

At 6 months: Hep B, Polio. These contain 62.5 mcg mercury which is 78 times above the 'safe' level.

At 15 months: The child receives another 50 mcg mercury which is 41 times above the 'safe' level.

Low levels of mercury during critical stages of development have been associated with neurological disorders in children including ADD learning difficulties, autism and speech delays. Wonder why we have an epidemic of these conditions? As of 2001, up to twenty-four vaccines are recommended from birth to eighteen years.

Reprinted from Sistah Qaraandin's M.O.T. Healthzine Issue #1